

Mitefauna (Arachnida: Acari) associated to grapevine, *Vitis vinifera* L. (Vitaceae), in the municipalities of Bento Gonçalves and Candiota, Rio Grande do Sul, Brazil

Crisna Letícia Klock¹, Liana Johann², Marcos Botton³ and Noeli Juarez Ferla^{1*}

- 1 Centro Universitário Univates, Museu de Ciências Naturais, Laboratório de Acarologia. Avelino Talini, 171. CEP 95900-000. Lajeado, RS, Brasil.
 - 2 Pontifícia Universidade Católica do Rio Grande do Sul, Programa de Pós-Graduação em Zoologia. Ipiranga, 6681. CEP 90619-900. Porto Alegre, RS, Brasil.
 - 3 Embrapa Uva e Vinho. Livramento, 515. CEP 95700-000. Bento Gonçalves, RS, Brasil.
- * Corresponding author. E-mail: niferla@univates.br

ABSTRACT: The mitefauna associated to Merlot and Chardonnay grapevine cultivars and associated plants in the municipalities of Bento Gonçalves and Candiota, Rio Grande do Sul was investigated. The study was developed between October 2006 and September 2007, where 20 grapevine plants were randomly chosen from each municipality and monthly sampled. Three leaves of each plant were taken. A total of 11,598 mites belonging to 14 families and to 52 species were found. Fifty-nine percent of the total specimens were collected in Candiota, being 93% associated to the Merlot cultivar. Higher species richness was observed on associated plants. Phytoseiidae showed the highest species richness, with ten species, and Eriophyidae showed the highest abundance, with 8,675 specimens. *Calepitrimerus vitis* (Nalepa, 1905) and *Polyphagotarsonemus latus* (Banks, 1904) were the most common phytophagous mites, while *Neoseiulus californicus* (McGregor, 1954) and *Pronematus anconai* (Baker, 1943) were the most common predators.

INTRODUCTION

Grapevine, *Vitis vinifera* L. (Vitaceae), was introduced in Brazil at 1532 by Martin Afonso de Souza expedition to São Paulo state (Lorenzi *et al.* 2006). Although it is cultivated today in nearly all regions of the country, the so called 'Serra Gaucha' region of the state of Rio Grande do Sul has currently the greatest vineyards and wineries concentration. More recently, vineyards have been implanted in the well-drained sandy soils of the flatter state region bordering Uruguay and Argentina (Botton *et al.* 2003; Mello 2008), that seems to be the region with the greatest potential for wine quality.

Grapevine plants are attacked by diseases, insects and mites. Amongst the phytophagous mites, the most important are those belonging to Eriophyidae, Tarsonemidae, Tenuipalpidae and Tetranychidae, since they frequently reach damage level in the vineyards (Reis and Melo 1984; Johann *et al.* 2009; Schruft 1985; Monteiro 1994; Duso and De Lillo 1996; Botton *et al.* 2003; Ferreira *et al.* 2006).

The most important eriophyid mites associated to vineyards are *Calepitrimerus vitis* (Nalepa, 1905) and *Colomerus vitis* (Pagenstecher, 1857) (Duso and De Lillo 1996; Jeppson *et al.* 1975). Three reported strains of *Colomerus vitis*, being each one characterized by the type of injury produced, are: the bud strain, the erineum strain and leaf-curling strain (Jeppson *et al.* 1975; Flechtman 1979; Duso and De Lillo 1996). The symptoms caused by *Calepitrimerus vitis* was brown scarification and necrosis on the inside of outer bud scales may be found under these mites attack, causing death of the growing point and sometimes of entire buds before sprouting. Injuries caused on shoots are drying, shortened internodes and development of lateral shoots. Injuries on leaves depend

on the season and leaf age. Infestation on growing leaves causes deformation due to irregular unfolding of the leaf tissues, with lacerations of the leaf margin, chlorotic spots often with necrosis in the center, and premature leaf fall (Duso and De Lillo 1996). *Calepitrimerus vitis* population was already observed on Alfrocheiro, Cabernet Sauvignon and Pinot Noir cultivars in Candiota, Encruzilhada do Sul and Bento Gonçalves regions, Rio Grande do Sul (Ferla *personal observations*; Johann *et al.* 2009).

Polyphagotarsonemus latus (Banks, 1904) (Tarsonemidae) develops mainly on young leaves, turning down their edges, causing browning and necrosis (Monteiro 1994). This species was already reported in vineyards in the state of Rio Grande do Sul (Johann *et al.* 2009), however it has not been considered of economic importance so far (Botton *et al.* 2005).

Several Tetranychidae species are associated to grapevine (Bolland *et al.* 1998), being *Panonychus ulmi* (Koch, 1836) the most important species in Northern Europe. *Panonychus ulmi* feeds mainly on the underside of leaves and its attack is particularly harmful in early spring and late summer. In summer, feeding of the mites causes bronzing of the leaves and affects the quality of fruits (Schruft, 1985). In Brazil, it was recently associated to viticulture (Ferla and Botton 2008). The attack diminishes the plant's photosynthetic surface, reduces the branches growing, the size and the color of the fruits and may cause premature leaf fall, interfering on the flowering and fructification in the following year (McMurtry *et al.* 1970; Croft 1975; Schruft 1985; Lorenzato 1987; Sousa 1996). *Oligonychus mangiferus* (Rahman and Punjab, 1940) and *Allonychus brasiliensis* (McGregor, 1950) were also reported from Brazil (Johann *et al.* 2009).

Although predatory mites from different families have

already been associated to grapevine, few studies about their predation capacity were carried out. Phytoseiid mites are efficient natural enemies of pest mites of several crops, and its presence is frequently associated with eriophyid mites in Rio Grande do Sul (Ferla *pers. obs.*), with *Euseius alatus* DeLeon, 1966, *Euseius brazilli* (ElBanhawy, 1975), *Neoseiulus fallacis* (Garman, 1948) and *Neoseiulus tunus* Denmark and Muma, 1967 already reported on grapevine (Johann *et al.* 2009; Monteiro 1994).

Despite the economic importance and the damage caused by some phytophagous mites on grapevine in Brazil, little is known about the species associated to this culture. However, because of the increase of the production area and the expansion in the border region of Uruguay and Argentina, it is fundamental to know the mitefauna associated and the damage caused by the most abundant and frequent species. Only then, a management plan to control mites in this culture can be suggested.

The present work aimed to identify the mitefauna associated to Chardonnay e Merlot varieties and associated plants during the harvest of 2006/2007 in the municipalities of Bento Gonçalves and Candiota, in the state of Rio Grande do Sul, Brazil.

MATERIALS AND METHODS

This study was carried out in vineyards of Chardonnay and Merlot varieties located in the municipalities of Bento Gonçalves (29°13' S, 51°33' W) and Candiota (31°28' S, 53°40' W), between October 2006 and September 2007 (Figure 1). Phytosanitary treatments with acaricides were not conducted during the study in the evaluated areas. The last treatment was applied on the harvest of 2005/2006.

The survey was conducted monthly with twenty plants of grapevine randomly selected from each area. A branch

from each plant was sampled, throughout the collection of three leaves from the apical, median and basal regions, respectively, totalizing 60 leaves per area. To evaluate the presence of mites on buds during senescence period, branches from twenty plants were removed and three buds by branch were sampled. Also, leaves from the five most common weeds around the grapevine rows were collected in enough quantity for one hour of sampling effort.

All collected leaves were introduced in paper bags, kept in a cooler box with Gelox® under low temperature ($\pm 15^{\circ}\text{C}$) and taken to the laboratory, where the material was stored under refrigeration for up to seven days until examination under a stereoscopic microscope for extraction of the mites. Mites found were collected with fine-tipped paintbrush and mounted with Hoyer's medium in microscopic slides (Jeppson *et al.* 1975). Slides were kept in a kiln during ten days at 50-60 °C to fix, to distend and to brighten up the specimens and to dry the medium. The identification was conducted under phase contrast microscope and with the help of the dichotomics keys (Amrine 1996; Baker *et al.* 1996; Baker and Tuttle 1994; Chant and McMurry 1994, 2003a and b, 2004a and b; 2005a, b and c; Halliday, Walter and Lindquist 1998; Krantz 2009; Lindquist 1986; Mantiolli, Ueckermann and Oliveira 2002; Smiley 1992; Summers and Price 1970). The plants species were identified by Dra. Elisete Maria de Freitas.

Voucher specimens of each species found were deposited on the mites reference collection of the Museu de Ciências Naturais of Centro Universitário UNIVATES (ZAUMCN), Lajeado, Rio Grande do Sul.

RESULTS AND DISCUSSION

A total of 11,598 mites belonging to 14 families were found. Fifty-nine percent of the specimens were collected in Candiota, being 93% associated to Merlot cultivar.

The mites found belonged to 52 species. Phytoseiidae showed the highest species richness, with ten species, followed by Tetranychidae and Stigmaeidae, with six species each, Tydeidae, with five species, and Ascidae, Eriophyidae and Tarsonemidae, with four species each. Eriophyidae showed the highest abundance, with 8,675 specimens, followed by Tarsonemidae, with 1,228 and Tydeidae, with 814.

The mite species and the host plants are presented below, along with the municipality, the cultivar, month and year of collection. Number of specimens is informed in brackets.

SUBORDER MESOSTIGMATA

ASCIDAE Voigts and Oudemans, 1905

Asca sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: *Plantago tomentosa* Lam. (1791) , VIII-2007 (1♀). **CANDIOTA**: **Chardonnay**: IV-2007 (1♀), VI-2007 (1♀), from *P. tomentosa*; **Merlot**: III-2007 (1♀), from *Solanum americanum* Mill. (1768).

Diseius sp.

Locality of the specimens examined: Brazil, Rio Grande




FIGURE 1. Location of the municipalities of Candiota and Bento Gonçalves, in the state of Rio Grande do Sul, Brazil.

do Sul, **BENTO GONÇALVES: Chardonnay:** VIII-2007 (1♀), from *P. tomentosa*.

Lasioseius sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** II-2007 (1♀), from *Taraxacum officinale* F. H. Wigg. (1780).

Leioseius sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** VIII-2007 (1♀), from *P. tomentosa*. **CANDIOTA: Chardonnay:** VI-2007 (1♀), from *P. tomentosa*.

PHYTOSEIIDAE Berlese, 1913

Amblyseius vitis Ferla and Silva, 2009

Amblyseius vitis Ferla and Silva, 2009: 509-510.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** VII-2007 (1♀), from *Stachys arvensis* L. (1763) ; IX-2007 (1♀), from *Rumex* sp.. **CANDIOTA: Chardonnay:** IV-2007 (1♀), from *P. tomentosa*.

Three adult females measurements: dorsal shield 443 (435-450) long and 304 (300-308) wide, j1 32 (25-37), j3 36 (32-38), j4 5 , j5 5, j6 5, J2 7 (7-8), J5 10 (7-13), z2 5, z4 6 (5-7), z5 6 (5-7), Z1 7 (5-8), Z4 214 (212-215), Z5 429 (405-442), s4 186 (175-195), S2 11 (10-12), S4 11 (10-12), S5 9 (8-10), r3 12 (12-13), R1 8 (7-10), SgI 71 (70-75), SgII 48 (45-50), SgIII 99 (97-100), StiIII 76 (75-80), SgIV 225 (223-227), StiIV 178 (173-185), StIV 145 (143-147), ST1-ST3 76 (75-80), ST2-ST2 91 (88-97), G-G 91 (87-97), ventrianal shield 135 (130-137) anterior wide, 101 (97-105) posterior wide and 105 (90-117) long, cervix of spermatheca 22 (20-25), fixed digit 44 (42-46), mobile digit 41 (38-43).

Previous records for Brazil: Rio Grande do Sul (Ferla and Silva 2009).

Comments: This species was described recently from the specimens collected on associated plants in vineyards in the state of Rio Grande do Sul.

Arrenoseius gaucho Ferla, Silva and Moraes, 2010

Arrenoseius gaucho Ferla, Silva and Moraes, 2010: 15-17

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** VIII-2007 (3♀1♂), from *P. tomentosa*; II-2007 (2♀), from *Solanum mauritianum* Scopoli (1788) **Merlot:** VIII-2007 (1♀), from *Plantago lanceolata* L. (1753); X-2006 (4♀3♂), XI-2006 (2♀), from *P. tomentosa*; V-2007 (2♀), from *Synedrella nodiflora* (L.) Gaertn (1791). **CANDIOTA: Chardonnay:** IV-2007 (3♀2♂), VI-2007 (2♀), from *P. tomentosa*; VI-2007 (2♀), from *Rumex* sp.. **Merlot:** VI-2007 (1♀), from *Blainvillea* sp.; I-2007 (1♀1♂), III-2007 (1♀), from *Richardia brasiliensis* Gomes (1801); VI-2007 (1♀), from *Senecio* sp..

Ten adult females measurements: dorsal shield 308 (295-325) long and 223 (207-245) wide, j1 21 (18-25), j3 20 (15-35), j4 11 (10-13), j5 9 (5-13), j6 10 (5-13), J2 11 (10-13), J5 9 (8-10), z2 19 (13-30), z4 12 (8-15), z5 7 (5-8), Z1 13 (8-15), Z4 66 (53-73), Z5 64 (58-70), s4 47

(30-65), S2 24 (13-28) , S4 11 (8-13), S5 10 (8-13), r3 20 (15-25), R1 13 (10-15), SgIV 46 (38-55), StiIV 30 (18-38) , StIV 41 (38-58), ST1-ST3 57 (53-63), ST2-ST2 68 (65-73), G-G 69 (65-78), ventrianal shield 98 (90-103) anterior wide, 83 (75-88) posterior wide and 103 (95-113) long, spermatheca cervix 16 (10-28), fixed digit 33 (31-36), mobile digit 30 (28-33).

Previous records for Brazil: Rio Grande do Sul (Ferla et al. 2010).

Euseius ho (De Leon, 1965)

Amblyseius (Euseius) ho DeLeon, 1965: 125.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** III-2007 (4♀), from *V. vinifera* leaf. **Merlot:** VII-2007 (1♀), from *P. tomentosa*.

Three adult females measurements: dorsal shield 343 (320-364) long and 252 (195-335) wide, j1 28 (27-30), j3 28 (23-38), j4 11 (7-18), j5 10 (8-10), j6 11 (10-13), J2 13 (12-13), J5 6 (5-8), z2 14 (12-15), z4 17 (15-18), z5 9 (8-10), Z1 13 (12-13), Z4 13 (12-15), Z5 52 (50-53), s4 28 (25-30), S2 15 (13-15), S4 17 (15-18), S5 18 (15-20), r3 13 (12-15), R1 11 (10-13), SgI 24 (23-25), SgII 24 (23-25), SgIII 27 (25-28), StiIII 23 (18-25), SgIV 33 (30-38), StiIV 27 (25-30), StIV 49 (45-52), ST1-ST3 59 (55-60), ST2-ST2 67 (65-70), G-G 75 (70-80), ventrianal shield 56 (50-62) anterior wide, 69 (57-75) posterior wide and 105 (100-105) long, spermatheca cervix 23 (22-23), fixed digit 24 (20-25), mobile digit 22 (20-25).

Previous records for Brazil: Bahia, Ceará, Paraíba, Pernambuco, Rio Grande do Sul, Santa Catarina and São Paulo (Denmark and Muma 1973; Moraes and McMurtry 1983; Moraes et al. 1990; Gondim Jr. and Moraes 2001; Zacarias and Moraes 2001; Ferla et al. 2005).

Comments: This species was reported in citrus and yerba mate crops in Taquari Valley, Rio Grande do Sul (Ferla et al. 2005). Species of this genus are common in native vegetation in that state (Ferla and Moraes 2002). *Euseius* species are polyphagous, but the reproductive potential of many species is highest on pollen (McMurtry and Croft 1997).

Euseius inouei (Ehara and Moraes, 1998)

Amblyseius (Euseius) inouei Ehara and Moraes, 1998: 59.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** V-2007 (1♀), from *Rumex* sp.; IV-2007 (1♀1♂), from *V. vinifera* leaf; V-2007 (2♀), VI-2007 (1♀), from *V. vinifera* bud.

Four adult females measurements: dorsal shield 338 (318-350) long and 210 (205-220) wide, j1 28 (28-30), j3 24 (12-30), j4 13 (10-17), j5 16 (12-15), j6 17 (15-18), J2 17 (15-22), J5 7 (6-8), z2 19 (17-23), z4 22 (20-25), z5 16 (15-18), Z1 17 (12-20), Z4 22 (17-25), Z5 61 (57-65), s4 31 (27-35), S2 22 (20-25), S4 27 (23-30), S5 29 (25-32), r3 15 (13-18), R1 14 (12-15), SgI 20 (18-25), SgII 21 (20-23), SgIII 27 (25-30), StiIII 21 (17-23), SgIV 39 (37-48), StiIV 24 (18-28), StIV 47 (43-50), ST1-ST3 60 (55-72), ST2-ST2 68 (63-70), G-G 76 (73-80), ventrianal shield 63 (57-75) anterior wide, 77 (74-87) posterior wide and 99 (95-103) long, spermatheca cervix 26 (21-28), fixed digit 25 (23-25), mobile digit 22 (20-23).

Previous records for Brazil: Rio Grande do Sul (Ferla and Moraes 2002).

Comments: The first record of this species in Brazil is by Ferla and Moraes (2002) in native and cultivated plants of Rio Grande do Sul.

Neoseiulus anonymus (Chant and Baker, 1965)

Amblyseius anonymus Chant and Baker, 1965:21

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay**: XI-2006 (1♀), from *V. vinifera* leaf.

The adult female measurement: dorsal shield 363 long and 180 wide, j1 20, j3 38, j4 28, j5 28, j6 40, J2 50, J5 10, z2 18, z4 38, z5 28, Z1 45, Z4 65, Z5 78, s4 50, S2 63, S4 35, S5 35, r3 33, R1 30, StIV 50, ST1-ST3 58, ST2-ST2 63, G-G 65, ventrianal shield 85 anterior wide and 75 posterior wide, spermatheca cervix 11, fixed digit 30, mobile digit 25.

Previous records for Brazil: Alagoas, Bahia, Paraíba, Paraná, Pernambuco, Rio Grande do Sul, Santa Catarina and São Paulo (Moraes et al. 2004).

Comments: This species belongs to a group of predators that show high oviposition rate when feeding on tetranychid mites (Ferla and Moraes 2003).

Neoseiulus californicus (McGregor, 1954)

Typhlodromus californicus McGregor, 1954: 89.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay**: XI-2006 (1♀), I-2007 (2), II-2007 (2♀1♂), from *V. vinifera* leaf. **Merlot**: V-2007 (1♀), from *Bidens pilosa* L. (1753); II-2007 (1♀), from *Conium maculatum* L. (1753); IV-2007 (1♀), from *Sonchus oleraceus* L. (1753); I-2007 (9♀5♂), II-2007 (10♀2♂), IV-2007 (6♀), from *V. vinifera* leaf; V-2007 (2♀), from *V. vinifera* bud. **CANDIOTA: Chardonnay**: VIII-2007 (1♀), from *Conyza canadensis* (L.) Cronquist (1943); IV-2007 (1♀), from *P. tomentosa*; IV-2007 (1♀), from *S. sataremnensis*; VI-2007 (1♀), from *S. chilensis*; I-2007 Meyen (1834) (5♀3♂), II-2007 (9♀3♂), III-2007 (23♀10♂), IV-2007 (35♀7♂), from *V. vinifera* leaf. **Merlot**: IV-2007 (1♀), from *B. pilosa*; VI-2007 (1♀), from *Blainvillea* sp.; VI-2007 (4♀1♂), from *Senecio* sp.; XI-2006 (1♀), I-2007 (23♀6♂), II-2007 (5♀), III-2007 (11♀2♂), IV-2007 (6♀), from *V. vinifera* leaf; V-2007 (3♀), from *V. vinifera* bud.

Seven adult females measurements: dorsal shield 352 (325-367) long and 168 (157-175) wide, j1 19 (17-22), j3 25 (22-32), j4 17 (12-22), j5 17 (15-22), j6 21 (17-25), J2 27 (22-30), J5 12 (10-12), z2 23 (17-25), z4 27 (20-32), z5 19 (17-20), Z1 26 (22-30), Z4 46 (42-50), Z5 64 (62-65), s4 28 (22-35), S2 34 (30 - 40), S4 31 (27-35), S5 27 (25-30), r3 21 (17-25), R1 20 (17-22), StIV 46 (40-52), ST1-ST3 60 (50 -65), ST2-ST2 60 (55-65), G-G 61 (55-68), ventrianal shield 107 (95-110) anterior wide, 87 (80-97) posterior wide and 116 (105-132) long, spermatheca cervix 12 (10-15), fixed digit 28 (25-32), mobile digit 24 (20-27).

Previous records for Brazil: Rio Grande do Sul (Ferla and Moraes 1998, 2002).

Comments: Specialized on tetranychid mites control (McMurtry and Croft 1997). It has been released on apple crop to control *P. ulmi* and *Tetranychus urticae* (Koch 1836) in Rio Grande do Sul and Santa Catarina (Monteiro 2002).

Neoseiulus tunus (DeLeon, 1967)

Typhlodromips tunus DeLeon, 1967: 29.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay**: III-2007 (2♀), from *V. vinifera* leaf.

Two adult females measurements: dorsal shield 301 (295-307) long and 179 (175-182) wide, j1 25, j3 26 (25-27), j4 9 (7-10), j5 9 (7-10), j6 15, J2 15, J5 8 (7-8), z2 21 (20-22), z4 32 (30-33), z5 12 (10-13), Z1 24 (23 -24), Z4 47 (45-48), Z5 75, s4 44 (40-47), S2 35 (33-37), S4 16 (15-17), S5 13 (12-13), r3 26 (25-27), R1 15, SgIV 18 (17-18), StIV 26 (25-27), ST1-ST3 64 (57-70), ST2-ST2 68 (67-68), GG 60 (58 -62), ventrianal shield 77 (75-78) anterior wide, 63 (60-65) posterior wide and 104 (100-107) long, spermatheca cervix 11 (10-12), fixed digit 32 (30 -33), mobile digit 29 (27-30).

Previous records for Brazil - Rio Grande do Sul, Santa Catarina and São Paulo (Moraes et al. 2004; Feres and Moraes 1998; Ferla and Moraes 1998, 2002; Lorenzato and Secchi 1993).

Comments: Species found in untreated apple orchards and native plants grown in the state of Rio Grande do Sul (Ferla and Moraes 1998, 2002).

Typhlodromalus aripo (DeLeon, 1967)

Typhlodromalus aripo DeLeon, 1967: 21.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay**: IX-2007 (1♀), from *Senecio* sp..

The adult female measurement: dorsal shield 354 long and 197 wide, j1 31, j3 35, j4 11, j5 12, j6 15, J2 15, J5 8, z2 17, z4 29, z5 11, Z1 20, Z4 46, Z5 68, s4 43, S2 33, S4 23, S5 11, r3 19, R1 16, SgI 20, SgII 19, SgIII 24, StiIII 15, SgIV 44, StiIV 24, StIV 69, ST1-ST3 67, ST2-ST2 63, G-G 78, ventrianal shield 66 anterior wide, 70 posterior wide 118 long, spermatheca cervix 19, fixed digit 38, mobile digit 34.

Previous records for Brazil: Bahia, Ceará, Maranhão, Pernambuco, Piauí, Rio Grande do Sul and São Paulo (Moraes and McMurtry 1983; Ferla and Moraes 1998, 2002; Denmark and Muma 1973; Feres and Moraes 1998; Zacarias and Moraes 2001; Moraes et al. 2004).

Comments: This is the most common predatory mite in plants associated with strawberry in Rio Grande do Sul (Ferla et al. 2007).

Typhlodromina tropica (Chant, 1959)

Typhlodromus tropicus Chant, 1959: 54.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot**: IV-2007 (1♀), from *V. vinifera* leaf.

The adult female measurement: 370 dorsal shield length 255 width, j1 10, j3 37, j4 32, j5 30, j6 45, J2 45, J5 5, 27 z2, z3 42, z4 50, z5 35, Z4 65, Z5 60, s4 57, s6 62, 70 S2, S5 45, r3 25, R1 17, St1-St3 57, St2-St2 67, GG 75, ventrianal shield 105 long, 75 wide and 70 posterior wide, spermatheca cervix 25.

Previous records for Brazil: São Paulo and Rio Grande do Sul (Denmark and Muma 1973; Ferla and Moraes 2002).

Typhlodromus ornatus (Denmark and Muma, 1973)

Amblydromella ornata Denmark and Muma, 1973: 270.

Typhlodromus ornatus Gondim and Moraes, 2001: 25.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** IV-2007 (2♀), V-2007 (1♀1♂), from *V. vinifera* bud. **Merlot:** VI-2007 (2♀), from *V. vinifera* bud. **CANDIOTA: Chardonnay:** VI-2007 (1♀), from *P. tomentosa*; III-2007 (3♀), IV-2007 (1♀1♂), VI-2007 (1), from *V. vinifera* leaf. **Merlot:** VIII-2007 (2♀), from *V. vinifera* bud.

Seven adult females measurements: dorsal shield 357 (295-378) long and 187 (163-200) wide, j1 21 (12-25), j3 18 (13-21), j4 11 (8-13), j5 10 (10-15), j6 13 (10-18), J2 16 (13-20), J5 12 (9-15), z2 14 (10-18), z3 15 (13-23), z4 16 (13-23), z5 12 (10-15), Z4 24 (18-33), Z5 38 (35-48), s4 18 (15-23), s6 20 (18-25), S2 21 (18-25), S4 23 (18-25), S5 23 (15-38), r3 18 (15-18), R1 19 (15-25), StIV 18 (15-25), StIV 25 (20-30), ST1-ST3 70 (50-78), ST2-ST2 61 (50-73), G-G 59 (54-63), ventrianal shield 88 (70-100) anterior wide, 79 (60-93) posterior wide and 117 (108-135) long, spermatheca cervix 20 (15-28), fixed digit 31 (15-38), mobile digit 29 (25-33).

Previous records for Brazil: São Paulo (Gondim Jr. and Moraes 2001).

SUBORDER PROSTIGMATA

CUNAXIDAE Thor, 1902

Neocunaxoides sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **CANDIOTA: Chardonnay:** IV-2007 (1♀), VI-2007 (7♀), from *P. tomentosa*.

ERIOPHYIDAE Nalepa, 1898

Aceria sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** I-2007 (40♀), from *S. mauritianum*. **CANDIOTA: Merlot:** IV-2007 (3♀), from *S. americanum*; XI-2006 (1♀), from *V. vinifera* leaf.

Calepitrimerus vitis (Nalepa, 1905)

Epitrimerus vitis Nalepa, 1905: 268.

Previous assignment: *Phyllocoptes*, *Epitrimerus*, *Calepitrimerus* (Amrine and Stasny 1994).

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** XII-2006 (1), I-2007 (38♀2♂), II-2007 (4♀), III-2007 (1♀), from *V. vinifera* leaf; **Merlot:** XI-2006 (3♀), XII-2006 (8♀), I-2007 (2.423), II-2007 (15♀), III-2007 (6♀), IV-2007 (2♀), from *V. vinifera* leaf. **CANDIOTA: Chardonnay:** I-2007 (14♀), II-2007 (58♀), III-2007 (28♀4♂), IV-2007 (2♀), from *V. vinifera* leaf; **Merlot:** I-2007 (2♀), from *Sida* sp.; I-2007 (1♀), from *S. americanum*; X-2006 (13♀), XI-2006 (212♀21♂), XII-2006 (12♀), I-2007 (5.611), II-2007 (74♀), III-2007 (11♀), IV-2007 (2♀), from *V. vinifera* leaf.

Previous records for Brazil: Rio Grande do Sul (Ferla pers. obs.; D'Andrea 1951).

Comments: This species of eriophyid mite was reported on viniferous grapevine in Candiota, Rio Grande do Sul (Ferla pers. obs.).

Colomerus vitis (Pagenstecher, 1857)

Phytoptus vitis Pagenstecher, 1857: 51.

Previous assignment: *Phytoptus*, *Eriophyies*, *Calomerus* (Amrine and Stasny 1994).

Locality of the specimens examined: **BENTO GONÇALVES: Merlot:** XI-2006 (1♀), XII-2006 (1♀), from *V. vinifera* leaf; VI-2007 (1♀), IX-2007 (44♀), from *V. vinifera* bud. **CANDIOTA: Merlot:** IV-2007 (1♀), from *V. vinifera* leaf.

Previous records for Brazil: Rio Grande do Sul (Johann et al. 2009; Braga 1957).

Comments: High populations may attack leaves, producing galls, and buds, causing deformation or even death (Duso and De Lillo 1996).

Criotacus sidae Keifer, 1977

Criotacus sidae Keifer, 1977, C13: 3, pl. 2.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** V-2007 (14♀), from *Sida santaremnensis* L. (1753).

Comments: First record in Brazil.

IOLINIDAE Pritchard, 1956

Homeopronematus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** II-2007 (1♀), from *S. mauritianum*; I-2007 (3♀), from *V. vinifera* leaf.

Metapronematus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** XII-2006 (2♀), I-2007 (2♀), from *V. vinifera* leaf. **Merlot:** XI-2006 (1♀), from *V. vinifera* leaf.

Pronematus anconai Baker, (1943) 1944

Pronematus anconai Baker, (1943) 1944: 188-189.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** XII-2006 (2♀), IV-2007 (6♀1♂), from *P. tomentosa*; VII-2007 (2♀), from *Rumex* sp.; I-2007 (2♀), II-2007 (2♀), from *S. mauritianum*; V-2007 (25♀2♂), VI-2007 (6♀), from *Sonchus oleraceus* L. (1753); I-2007 (1♀), VII-2007 (1♀), from *T. officinale*; XII-2006 (19♀), I-2007 (85♀12♂), II-2007 (9♀), III-2007 (8♀), from *V. vinifera* leaf; VI-2007 (8♀), from *V. vinifera* bud. **Merlot:** XII-2006 (2♀), from *Paspalum* sp.; VI-2007 (5♀), from *Poa annua* L. (1753); I-2007 (11♀), IV-2007 (1♀), from *S. oleraceus*; I-2007 (1♀), from *Trifolium repens* L. (1753); I-2007 (30♀14♂), from *V. vinifera* leaf. **CANDIOTA: Chardonnay:** V-2007 (1♀), from *S. americanum*; II-2007 (1♀), IV-2007 (3), from *S. oleraceus*. **Merlot:** IV-2007 (1♀), from *Bidens pilosa* L.; I-2007 (1♀), from *Digitaria* sp.; IV-2007 (1♀), V-2007 (2♀), from *R. brasiliensis*; VII-2007 (1♀), from *Senecio selloi* Spreng. (1837); IV-2007 (1♀), from *S. americanum*.

Distribution: It was described from Mexico specimens (Baker 1968).

Comments: This species is an important alternative prey for the predatory mite of *Metaseiulus occidentalis* (Nesbitt) in California vineyards (Flaherty and Hoy 1971). It feeds primarily on pollen, the predation rate on spider mite eggs is low and it could not reproduce on leaves having only spider mites eggs (Knop and Hoy 1983).

PYEMOTIDAE Oudemans, 1937

Pyemotes sp.

Specimen examined: **BENTO GONÇALVES**: Brazil, Rio Grande do Sul, **Chardonnay**: VII-2007 (1♀), from *S. arvensis*.

Pygmephorus sp.

Locality of the specimens examined: **BENTO GONÇALVES**: **Chardonnay**: VIII-2007 (1♀), from *P. tomentosa*.

STIGMAEIDAE Oudemans, 1931

Agistemus sp.1

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: III-2007 (2♀), from *V. vinifera* leaf.

Agistemus sp.2

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Merlot**: IV-2007 (2♀), from *Galinsoga* sp..

Agistemus sp.3

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Merlot**: V-2007 (2♀), from *Sida santaremnensis* H. Monteiro.

Agistemus brasiliensis Matioli, Ueckermann and Oliveira, 2002

Agistemus brasiliensis Matioli, Ueckermann and Oliveira, 2002: 206.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: I-2007 (2♀1♂), from *V. vinifera* leaf. **Merlot**: XI-2006 (1♀), IV-2007 (1♀), from *V. vinifera* leaf.

Previous records for Brazil: Rio Grande do Sul and São Paulo (Matioli et al. 2002; Ferla et al. 2005).

Comments: Species associated with yerba mate in Rio Grande do Sul, with higher populations coinciding with eriophyid mites populations peaks (Ferla et al. 2005).

Agistemus floridanus Gonzalez, 1965

Agistemus floridanus Gonzalez, 1965: 42.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: XII-2006 (3♀), I-2007 (14♀6♂), II-2007 (25♀13♂), III-2007 (10♀), from *V. vinifera* leaf. **Merlot**: IV-2007 (1♀), from *P. tomentosa*; I-2007 (2♂), IV-2007 (27♀), from *V. vinifera* leaf; VI-2007 (1♀), from *V. vinifera* bud. **CANDIOTA**: **Merlot**: V-2007 (1♀), from *B. pilosa*; II-2007 (1♀), III-2007 (1♀), IV-2007 (2♂), from *V. vinifera* leaf.

Previous records for Brazil: Rio Grande do Sul and Mato Grosso (Ferla and Moraes 2003; Ferla et al. 2007).

Comments: Species reported in strawberry culture in Rio Grande do Sul (Ferla et al. 2007). It shows high oviposition levels when fed on tetranychid, eriophyid mites and pollen of *Typha angustifolia* L. (1753) (Ferla and Moraes 2003).

Zetzellia malvinae Matioli, Ueckermann and Oliveira, 2002

Zetzellia malvinae Matioli, Ueckermann and Oliveira,

2002: 111.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **CANDIOTA**: **Merlot**: VI-2007 (1♀), from *Senecio* sp.; VII-2007 (2♀), from *S. selloi*; II-2007 (1♀), IV-2007 (5), from *V. vinifera* leaf; V-2007 (1♀), VI-2007 (1♀), from *V. vinifera* bud.

Previous records in Brazil: São Paulo (Matioli et al. 2002).

Comments: It was described from specimens of citrus (Matioli et al. 2002).

TARSONEMIDAE Kramer, 1877

Acaronemus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: VIII-2007 (1♀), from *P. tomentosa*; IV-2007 (1♀), from *V. vinifera* bud. **Merlot**: VII-2007 (1♀), from *P. tomentosa*; V-2007 (1♀), from *Rumex* sp.; X-2006 (1♀), from *Trifolium Repens* L. (1753); VI-2007 (1♀), VIII-2007 (1♀), from *V. vinifera* bud. **CANDIOTA**: **Chardonnay**: X-2006 (1♀), from *Solanum* sp.; I-2007 (1♀), from *S. americanum*; I-2007 (1♀), from *V. vinifera* leaf; V-2007 (1♀), VII-2007 (1♀), from *V. vinifera* bud. **Merlot**: IV-2007 (1♂), from *B. pilosa*; V-2007 (1♂), from *R. brasiliensis*; V-2007 (1♀), from *Sida* sp.; II-2007 (1♀), IV-2007 (3♀1♂), from *V. vinifera* leaf; V-2007 (1♀), from *V. vinifera* bud.

Polyphagotarsonemus latus (Banks, 1904)

Tarsonemus latus Banks, 1904: 1553.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: I-2007 (5♀2♂), from *B. pilosa*; IV-2007 (1♀), from *P. tomentosa*; I-2007 (24♀9♂), from *S. mauritianum*; XII-2006 (2♀), I-2007 (656♀71♂), II-2007 (34♀), from *V. vinifera* leaf. **Merlot**: *Conium maculatum* L., I-2007 (1♀); I-2007 (84♀35♂), from *V. vinifera* leaf.

Previous records for Brazil: Pernambuco and Rio Grande do Sul (Ferreira et al. 2006; Johann et al. 2009; Soria et al. 1993)

Comments: As a grapevine pest, it causes branch shortening (Botton et al. 2005). In situations of high infestation, the leaves get coriaceous and brittle, and leaf fall may occur (Haji et al. 2001).

Tarsonemus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES**: **Chardonnay**: I-2007 (1♀), from *S. mauritianum*; VII-2007 (1♀), from *P. tomentosa*; I-2007 (1♀), III-2007 (1♀), from *V. vinifera* leaf; IV-2007 (6♀), V-2007 (31), VI-2007 (2♀), VII-2007 (2♀1♂), VIII-2007 (5♀), IX-2007 (3♀1♂), from *V. vinifera* bud. **Merlot**: VIII-2007 (1♀), from *P. lanceolata*; VII-2007 (1♀), VIII-2007(11), from *P. tomentosa*; VII-2007 (1♀), from *Rumex* sp.; XII-2006 (1♀), IV-2007 (1♀), from *V. vinifera* leaf; V-2007 (9♀), VI-2007 (27♀), VIII-2007 (2♀), from *V. vinifera* bud. **CANDIOTA**: **Chardonnay**: VI-2007(1♀), from *Baccharis trimera* Less. (1836); VIII-2007 (2♀), from *Conysa canadensis* L.; II-2007(1♀), IV-2007 (44♀8♂), from *V. vinifera* leaf; V-2007 (1♀), VI-2007 (1♀), VII-2007 (3♀), VIII-2007 (1♀), from *V. vinifera* bud. **Merlot**: IV-2007 (5♀), from *B. pilosa*; IX-2007 (2♀), from *Eupatorium buniifolium*

Hook and Arn. (1835); V-2007 (1♀), from *Galinsoga quadriradiata* Ruiz and Pav. (1794); IX-2007 (1♀), from *Gnaphalium spicatum* Lam. (1788); VIII-2007 (4♀), from *Senecio* sp.; V-2007 (1♀), from *Sida* sp.; IV-2007 (4♀), from *S. americanum*; III-2007 (2♀), IV-2007 (30♀7♂), from *V. vinifera* leaf; V-2007 (13♀), VI-2007 (2♀), VIII-2007 (21♀), IX-2007 (3♀), from *V. vinifera* bud.

Xenotarsonemus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** VIII-2007 (1♀), *P. lanceolata*; XI-2006 (1♀), VIII-2007 (1♀), from *P. tomentosa*. **CANDIOTA: Chardonnay:** VI-2007 (1♀), from *P. tomentosa*. **Merlot:** VII-2007 (2♂), IX-2007 (3♀), from *B. trimera*; IX-2007 (1♀), from *E. buniifalium*; IX-2007 (1♀), from *G. spicatum*; VI-2007 (1♀), from *Oxalis corniculata* L. (1753); IV-2007 (2♀), from *R. brasiliensis*.

TENUIPALPIDAE Berlese, 1913

Brevipalpus phoenicis (Geijskes, 1939)

Tenuipalpus phoenicis Geijskes, 1939: 23.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** IV-2007 (1♀), from *V. vinifera* leaf. **CANDIOTA: Merlot:** VII-2007 (1♀), IX-2007 (5♀), from *B. trimera*; VIII-2007 (18♀2♂), from *Gnaphalium* sp.; VIII-2007 (1♀), from *Rumex* sp..

Previous records for Brazil: São Paulo, Santa Catarina, Pernambuco and Ceará (Oliveira 1986; Lorenzato 1987; Ferla et al. 2005).

Comments: Reported as citrus leprosis mite, it is the vector of the CiLV virus, agent of the disease in citrus (Chiavegato 1987). It is also associated with coffee ringspot, where it transmits the CoRSV virus (Chagas 1973). It is not referred to cause economic damage on grapevine culture (Flechtmann 1979).

TETRANYCHIDAE Donnadieu, 1875

Mononychellus planki (McGregor, 1950)

Tetranychus planki McGregor, 1950: 300.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** V-2007 (1♀), from *S. santaremnensis*.

Previous records for Brazil: Broad distribution in Brazil (Moraes and Flechtmann 2008).

Comments: It is not a species of economic importance, but it may be considered a potencial cotton pest. It was found in soybean, *Glycine max* (L.) (1917) Merr., in Rio Grande do Sul (Roggia et al. 2008), where its populations seemed to be increasing in the latest years.

Oligonychus yothersi (McGregor, 1914)

Tetranychus yothersi McGregor, 1914: 355.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** X-2006 (1♀), from *V. vinifera* leaf. **Merlot:** X-2006 (1♂), from *V. vinifera* leaf.

Previous records for Brazil - Paraná, Rio Grande do Sul, Santa Catarina and São Paulo (Ferla et al. 2005).

Comments: It is known as yerba mate red mite. It attacks new and ripe leaves, causing brazing and defoliation in

severe attacks. During dry and warm periods it attacks in outbreak, usually disappearing after intense rainfall (Flechtmann 1979; Coll and Saini 1992).

Oligonychus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** I-2007 (1♀), from *T. officinale*.

Panonychus ulmi (Koch, 1836)

Tetranychus ulmi Koch, 1836: 11.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** I-2007 (10♀), III-2007 (2♂), from *V. vinifera* leaf. **Merlot:** X-2006 (2♀), XI-2006 (2♀2♂), I-2007 (24♀3♂), II-2007 (7♀), III-2007 (8♀), from *V. vinifera* leaf.

Previous records for Brazil: Rio Grande do Sul (Ferla and Botton 2008).

Comments: This species is also considered an apple pest, causing leaf brownish, reducing branch growth, size, fruit color and the sugar content, and also causing premature leaf fall. It interferes on the following year's flowering and fructification (Monteiro 2002). It was observed on grapevine in Rio Grande do Sul, where the leaves showed brownish, with reddish branches on adaxial face and premature leaf fall (Ferla and Botton 2008).

Petrobia sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** VI-2007 (8♀), from *P. annua*.

Tetranychus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** I-2007 (6♀), II-2007 (1♀), from *Galinsoga parviflora* Cav. (1794).

TYDEIDAE Kramer, 1877

Lorryia formosa Cooreman, 1958

Lorryia formosa Cooreman, 1958: 6.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** V-2007 (1♀), from *S. santaremnensis*; IV-2007 (1♀), from *V. vinifera* leaf.

Previous records for Brazil: São Paulo (Feres et al. 2003).

Comments: This mite is rather common in Brazil and has been collected from a rather wide range of plants (Flechtmann 1973, 1987; Feres et al. 2003; Hernandes and Feres 2006). Populations of this mite may cause sclerotization of the foliar limb followed by scaling in the area where they are concentrated (Jeppson et al. 1975).

Lorryia sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Merlot:** I-2007 (1♀), IV-2007 (1♀), from *V. vinifera* leaf. **CANDIOTA: Merlot:** V-2007 (1♀), from *V. vinifera* bud.

Metatritophydeus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** IX-2007 (1♀),

from *Senecio* sp.; I-2007 (4♀), from *V. vinifera* leaf; V-2007 (1♀), VI-2007 (1♀), IX-2007 (1♀), from *V. vinifera* bud.

Paralorryia sp.

Specimen examined: Brazil, Rio Grande do Sul, **CANDIOTA: Merlot:** IV-2007 (1♀), from *R. brasiliensis*.

Tydeus sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** V-2007 (3♀), from *B. pilosa*; X-2006 (1♀), VII-2007 (1♀), from *C. maculatum*; IV-2007 (3♀), IX-2007 (1♀), from *P. tomentosa*; IX-2007 (1♀), from *Senecio* sp.; VI-2007 (1♀), from *S. oleraceus*; XII-2006 (6♀), I-2007 (1♀), II-2007 (3♀), III-2007 (21♀), from *V. vinifera* leaf; IV-2007 (4♀), V-2007 (46♀11♂), VI-2007 (60♀5♂), VII-2007 (8♀), VIII-2007 (3♀), IX-2007 (1♀), from *V. vinifera* bud. **Merlot:** VII-2007 (32♀5♂), VIII-2007 (14♀), from *P. tomentosa*; VII-2007 (4♀), from *Rumex* sp.; VI-2007 (7♀), from *S. nodiflora*; VI-2007 (1♀), from *T. repens*; XI-2006 (2♀), XII-2006 (1♀), III-2007 (1♀), IV-2007 (52♀), from *V. vinifera* leaf; V-2007 (8♀), VI-2007 (107♀11♂), VII-2007 (28♀), VIII-2007 (4♀), from *V. vinifera* bud. **CANDIOTA: Chardonnay:** VIII-2007 (6♀), from *C. canadensis*; VI-2007 (6♀), from *P. tomentosa*; VIII-2007 (2♀), from *Rumex* sp.; IX-2007 (6♀), from *Senecio brasiliensis* Less.; V-2007 (3♀), from *S. americanum*; VI-2007 (8♀), from *Solidago chilensis* Meyen.; IV-2007 (1♀), from *S. oleraceus*; II-2007 (1♀), III-2007 (1♀), from *V. vinifera* leaf; IV-2007 (28♀), V-2007 (9♀), VI-2007 (39♀), VII-2007 (3♀), VIII-2007 (3♀), from *V. vinifera* bud. **Merlot:** IV-2007 (2♀), V-2007 (4♀), *B. pilosa*; V-2007 (2♀), from *G. quadriradiata*; IV-2007 (2♀), V-2007 (13♀), from *R. brasiliensis*; VI-2007 (4♀), from *Senecio* sp.; VI-2007 (7♀), from *S. brasiliensis*; VII-2007 (2♀), from *S. selloi*; V-2007 (1♀), from *Sida* sp.; IV-2007 (17♀), from *S. americanum*; V-2007 (3♀), from *S. oleraceus*; III-2007 (1♀), IV-2007 (77♀7♂), from *V. vinifera* leaf; V-2007 (26♀), VI-2007 (45♀), VII-2007 (6♀), VIII-2007 (6♀), from *V. vinifera* bud.

SUBORDER ASTIGMATA

HISTIOSTOMIDAE Hughes, 1976

Histiostoma sp.

Specimen examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** VII-2007 (1♀), from *Rumex* sp. L..

WINTERSCHMIDTIIDAE Oudemans, 1923

Czenspinskia sp.

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** XII-2006 (1♀), from *V. vinifera* leaf. **Merlot:** VII-2007 (1♀), from *V. vinifera* bud.

SUBORDER ORIBATIDA

Locality of the specimens examined: Brazil, Rio Grande do Sul, **BENTO GONÇALVES: Chardonnay:** VIII-2007 (1♀), from *P. tomentosa*; VII-2007 (3♀), from *S. arvensis*; IX-2007

(3♀), from *V. vinifera* bud. **Merlot:** VIII-2007 (3♀), from *P. lanceolata*; X-2006 (2♀), IV-2007 (5♀), VI-2007 (2♀), VII-2007 (1♀1♂), VIII-2007 (4♀2♂), from *P. tomentosa*; V-2007 (1♀), from *Rumex* sp.; V-2007 (1♀), from *S. santaremnensis*. **CANDIOTA: Chardonnay:** VIII-2007 (1♀), from *B. trimera*; VIII-2007 (1♀), from *C. canadensis*; VII-2007 (3♀1♂), from *Gnaphalium* sp.; IV-2007 (1♀), VI-2007 (5♀), from *P. tomentosa*; VII-2007 (3♀), VIII-2007 (1♀), from *Rumex* sp.. **Merlot:** VI-2007 (3♀), from *Blainvillea* sp.; VII-2007 (3♀), from *Gnaphalium* sp.; VIII-2007 (1♀), from *Senecio* sp.

Mites found in each plant species associated with grapevine sampled are presented in Table 1. The grapevine culture presented great mite richness, higher number of species in Bento Gonçalves (46 species) than in Candiota (21 species). This fact may be related to the recent introduction of the culture in that region or to distinct environmental factors and climate, once Bento Gonçalves is located in the northeast upper slopes, where rainfall and humidity are high, while Candiota is located in west border of the state and has a drier climate and low relative humidity, mainly during the ripening period.

Calepitrimerus vitis and *Cololomerus vitis* were the most abundant species, with *Cal. vitis* being the most common species in the leaves, indicating it is probably the more important phytophagous mite in grapevine in the state. Their monitoring is necessary, because higher populations may cause bud's death decreasing the vegetative growth and leaves brownish with decrease of plant photosynthetic capacity (Duso and De Lillo 1996). The interference on plant photosynthetic capacity also interferes in the quality of the wine produced, since good wines are produced with grapes of good quality (Vencato 2007).

Colomerus vitis was noted only in Merlot cultivar in the two places sampled and populations are much higher in buds. This species may cause deformation in buds and new leaves or still cause the appearance of eriosis of the leaves (Jeppson et al. 1975; Flechtman 1979; Duso and De Lillo 1996).

Just two tetranychid mite species were observed in Bento Gonçalves. *Oligonychus yothersi* was found at low density and does not present economic importance to the culture. *Panonychus ulmi*, present in the two cultivars studied in Bento Gonçalves, was recently reported in the culture in Brazil, with damages described by Ferla and Botton (2008). Commonly associated to Europe grapevine (Jeppson et al. 1975; Schruft 1985; Bolland et al. 1998), it becomes important to the culture in Rio Grande do Sul. Although *P. ulmi* has not appeared in high density, its finding denotes that this species should be inserted in integrated pest management programs in the state of Rio Grande do Sul.

Four Tarsonemid species were identified and the most abundant were *Polyphagotarsonemus latus* and *Tarsonemus* sp.. *Polyphagotarsonemus latus* was present just in Bento Gonçalves and only on apical leaves. It occurred mainly in January on new leaves after pruning green, suggesting that species did not interfere in vineyard formation (Johann et al. 2009). Nothing is known about the damage caused by this species in that period in grapevine plants.

TABLE 1. Mitefauna of the uncultivated plants associated with grapevine Chardonnay (C) and Merlot (M), in the municipalities of Bento Gonçalves (BG) and Candiota (CA), Rio Grande do Sul, Brazil.

VEGETAL TAXON	MITES ASSOCIATED	BG		CA	
		C	M	C	M
AMARANTHACEAE					
<i>Amaranthus</i> sp.	-				
<i>Amaranthus deflexus</i> L. (1771)	-				
<i>Amaranthus hybridus</i> L. (1753)	-				
APIACEAE					
	<i>Neoseiulus californicus</i>		1		
<i>Conium maculatum</i>	<i>Polyphagotarsonemus latus</i>		1		
	<i>Tydeus</i> sp.	2			
ASTERACEAE					
	<i>Acaronemus</i> sp.				1
	<i>Agistemus floridanus</i>				1
	<i>Neoseiulus californicus</i>				1
<i>Bidens pilosa</i>	<i>Polyphagotarsonemus latus</i>	7			
	<i>Pronematus anconai</i>		1		1
	<i>Tarsonemus</i> sp.				5
	<i>Tydeus</i> sp.	3			6
<i>Baccharis trimera</i>	<i>Breviphalpus phoenicis</i>			10	6
	<i>Arrenoseius gauch</i> o				1
<i>Blainvillea</i> sp.	<i>Neoseiulus californicus</i>				1
	Oribatida				3
<i>Conyza</i> sp.	-				
	<i>Neoseiulus californicus</i>			1	
<i>Conysa canadensis</i>	<i>Tarsonemus</i> sp.			2	
	<i>Tydeus</i> sp.			6	
<i>Emilia sonchifolia</i> L. (1834)	-				
<i>Eupatorium buniifalium</i>	<i>Tarsonemus</i> sp.				2
	<i>Xenotersonemus</i> sp.				1
<i>Galinsoga</i> sp.	<i>Agistemus</i> sp. ²		2		
<i>Galinsoga parviflora</i>	<i>Tetranychus</i> sp.	7			
<i>Galinsosa quadriradiata</i>	<i>Tarsonemus</i> sp.				1
	<i>Tydeus</i> sp.				1
<i>Gnaphalium</i> sp.	<i>Breviphalpus phoenicis</i>				20
	Oribatida			4	3
<i>Ganphalium spicatum</i>	<i>Tarsonemus</i> sp.				1
	<i>Xenotersonemus</i> sp.				1
<i>Hypochoeris radicata</i> L. (1753)	-				
	<i>Arrenoseius gauch</i> o				1
	<i>Metetriophtydeus</i> sp.	1			
	<i>Neoseiulus californicus</i>				5
<i>Senecio</i> sp.	Oribatida				1
	<i>Tarsonemus</i> sp.				4
	<i>Tydeus</i> sp.	1			5
	<i>Typhlodromalus aripo</i>	1			
	<i>Zetzellia malvinae</i>				1
<i>Senecio brasiliensis</i>	<i>Tydeus</i> sp.			6	7
	<i>Pronematus anconai</i>				1
<i>Senecio selloi</i>	<i>Tydeus</i> sp.				2
	<i>Zetzellia malvinae</i>				2
<i>Solidago chilensis</i>	<i>Neoseiulus californicus</i>			1	
	<i>Tydeus</i> sp.			8	
	<i>Neoseiulus californicus</i>		1		
<i>Sonchus oleraceus</i>	<i>Pronematus anconai</i>	33	12	4	
	<i>Tydeus</i> sp.	1		1	3

TABLE 1. CONTINUED.

VEGETAL TAXON	MITES ASSOCIATED	BG		CA	
		C	M	C	M
<i>Synedrella nodiflora</i>	<i>Arrenoseius gaucho</i>		2		
	<i>Tydeus</i> sp.		7		
	<i>Lasioseius</i> sp.		1		
<i>Taraxacum officinale</i>	<i>Oligonychus</i> sp.	1			
	<i>Pronematus anconai</i>	2			
CARYOPHYLLACEAE					
<i>Stellaria media</i> (L.) Vill. (1784)	-				
GRAMINEAE					
<i>Poa annua</i>	<i>Petrobia</i> sp.		8		
	<i>Pronematus anconai</i>		5		
FABACEAE					
<i>Medicago hispida</i> Gaertn. (1791)	-				
<i>Medicago lupulina</i> L. (1753)	-				
<i>Trifolium pratenses</i> L. (1753)	-				
<i>Trifolium repens</i>	<i>Acaronemus</i> sp.		1		
	<i>Pronematus anconai</i>		1		
	<i>Tydeus</i> sp.		1		
<i>Vicia sativa</i> Guss. (1828).	-				
LAMIACEAE					
<i>Leonurus sibiricus</i> L. (1753)	-				
<i>Stachys arvensis</i>	<i>Amblyseius vitis</i>	1			
	Oribatida	3			
	<i>Pyemotes</i> sp.	1			
LYTHRACEAE					
<i>Cuphea</i> sp.	-				
MALVACEAE					
<i>Malvastrum coromandelianum</i> L.(1857)	-				
<i>Sida</i> sp.	<i>Acaronemus</i> sp.				1
	<i>Calepitrimerus vitis</i>				2
	<i>Tarsonemus</i> sp.				1
	<i>Tydeus</i> sp.				1
<i>Sida rhombifolia</i> L. (1753)	-				
<i>Sida santaremnensis</i>	<i>Agistemus</i> sp. ³		2		
	<i>Crioticus sidae</i>		14		
	<i>Lorryia formosa</i>		1		
	<i>Mononychellus planki</i>	1			
	<i>Neoseiulus californicus</i>			1	
	Oribatida		1		
<i>Sida spinosa</i> L. (1753)	-				
OXALIDACEAE					
<i>Oxalis corniculata</i>	<i>Xenotarsonemus</i> sp.				1
PLANTAGINACEAE					
<i>Plantago lanceolata</i>	<i>Arrenoseius gaucho</i>		1		
	Oribatida		3		
	<i>Tarsonemus</i> sp.		3		
	<i>Xenotarsonemus</i> sp.		1		

TABLE 1. CONTINUED.

VEGETAL TAXON	MITES ASSOCIATED	BG		CA	
		C	M	C	M
<i>Plantago tomentosa</i>	<i>Acaronemus</i> sp.	1	1		
	<i>Agistemus floridanus</i>		1		
	<i>Amblyseius vitis</i>			1	
	<i>Asca</i> sp.	2		1	
	<i>Diseius</i> sp.	1			
	<i>Euseius ho</i>		1		
	<i>Arrenoseius gaucha</i>	4	9	7	
	<i>Leioseius</i> sp.	1		1	
	<i>Neocunaxoides</i> sp.			8	
	<i>Neoseiulus californicus</i>			1	
	Oribatida	1	17	16	
	<i>Polyphagotarsonemus latus</i>	1			
	<i>Pronematus anconai</i>	9			
	<i>Pygmephorus</i> sp.	1			
	<i>Tarsonemus</i> sp.	1	12		
	<i>Tydeus</i> sp.	4	51	6	
	<i>Ttyphlodromus ornatus</i>			1	
	<i>Xenotarsonemus</i> sp.		2	1	
PLYGONACEAE					
<i>Rumex</i> sp.	<i>Acaronemus</i> sp.		1		
	<i>Amblyseius vitis</i>	1			
	<i>Breviphalpus phoenicis</i>				1
	<i>Euseius inouei</i>		1		
	<i>Arrenoseius gaucha</i>			1	
	<i>Histiostoma</i> sp.	1			
	Oribatida		1	4	
	<i>Pronematus anconai</i>	2			
	<i>Tarsonemus</i> sp.		1		
	<i>Tydeus</i> sp.		4	2	
POACEAE					
<i>Brachiaria</i> sp. (Trin.) Griseb. (1853)	-				
<i>Cenchrus</i> sp. L. (1753)	-				
<i>Digitaria</i> sp.	<i>Pronematus anconai</i>				1
	<i>Paspalum</i> sp.		2		
<i>Digitaria sanguinalis</i> (L.) Scop. (1771)	-				
<i>Eleusine distachya</i> Nees. (1829)	-				
<i>Eragrostis</i> sp. Wolf (1776)	-				
<i>Lolium multiflorum</i> Lam. (1779)	-				
<i>Paspalum</i> sp.	-				
PORTULACACEAE					
<i>Portulaca oleraceae</i> L. (1753)	-				
RUBIACEAE					
<i>Richardia brasiliensis</i>	<i>Acaronemus</i> sp.				1
	<i>Czenpinskia</i> sp.				1
	<i>Arrenoseius gaucha</i>				3
	<i>Paralorryia</i> sp.				1
	<i>Pronematus anconai</i>				3
	<i>Tydeus</i> sp.				15
	<i>Xenotarsonemus</i> sp.				2
<i>Spermacoce verticillata</i>	<i>Tetranychus</i> sp.			2	
SOLANACEAE					
<i>Nicotiana tabacum</i> L.(1753)	-				
<i>Solanum</i> sp.	<i>Acaronemus</i> sp.			1	

TABLE 1. CONTINUED.

VEGETAL TAXON	MITES ASSOCIATED	BG		CA	
		C	M	C	M
<i>Solanum americanum</i>	<i>Acaronemus</i> sp.			1	
	<i>Aceria</i> sp.				3
	<i>Asca</i> sp.				1
	<i>Brevipalpus phoenicis</i>			1	
	<i>Calepitrimerus vitis</i>				1
	<i>Pronematus anconai</i>			1	1
	<i>Tarsonemus</i> sp.				4
	<i>Tydeus</i> sp.			3	17
<i>Solanum mauritianum</i>	<i>Aceria</i> sp.	40			
	<i>Arrenoseius gaucho</i>	2			
	<i>Homeopronematus</i> sp.	1			
	<i>Polyphagotarsonemus latus</i>	33			
	<i>Pronematus anconai</i>	4			
	<i>Tarsonemus</i> sp.	1			
URTICACEAE					
<i>Parietaria debilis</i> G.Forst. (1786)	-				
TOTAL		176	174	103	149

Amongst the predatory mites, the Phytoseiidae presented the higher diversity, mainly in Bento Gonçalves. *Neoseiulus californicus*, the most common species, was observed in both localities, associated with *Cal. vitis*. The phytoseiid mite diversity in Serra Gaucha can be related to the presence of tetranychids mites, while low diversity was found in Candiota where this phytophagous group was not found. Tetranychid mites are favorable preys to Phytoseiidae predator mites and the presence of this preys can hold higher number of prey (McMurtry and Croft 1997; Moraes 2002). *Neoseiulus californicus* has been utilized in program of biologic control of *P. ulmi* in apple culture with relative success in the states of Rio Grande do Sul and Santa Catarina (Monteiro 2002). In Candiota, the occurrence of *N. californicus* was reported in the same period in which *Cal. vitis* was found, indicating possible interactions. According to Moraes (2002), there are some species of Eriophyidae that are quite favorable preys to development of phytoseiid mites. Johann *et al.* (2009), also observed high population of this predator associated with *Cal. vitis* in Candiota.

The Iolinidae was the most frequent predatory mite and *P. anconai* was the most abundant species associated with *Cal. vitis* and *P. latus* populations, suggesting that it can be an important natural enemy of these species. Species of this family have been reported by Laing and Knop (1982) and Perrin and McMurtry (1996) as predators of eriophyid mites. In Germany, *Pronematus staerki* Schruft is an important natural enemy of *Cal. vitis* (Duso and De Lillo 1996). *Pronematus anconai* was previously reported in the grapevine varieties Alfrocheiro, Cabernet Sauvignon and Pinot Noir in the cities of Bento Gonçalves and Candiota, in Rio Grande do Sul (Johann *et al.* 2009).

The Tydeidae also showed great species richness (five species) and *Tydeus* sp. was the most abundant both the localities evaluated. The species was more frequent on

leaves during the senescent period, in April and May, and in the bud beginning period. Species of this genus were reported as mites with different feeding habits by several authors. *Tydeus caudatus* Dugès, *T. goetzi* Schruft and *T. grabouwi* Meyer and Ryke were reported as natural enemies of *Cal. vitis* and *Col. vitis* in Europe and South Africa (Duso *et al.* 2004). Representatives of this family were also observed feeding on fungi (Hessein and Perring 1986). The tydeids may have high ecologic importance in agroecosystems to be considered as alternative food to phytoseiid mites (Strickler *et al.* 1987; McMurtry and Croft 1997; Ferla and Moraes 1998). The consumption of alternative food keeps the predators in the environment where the preferential prey is absent (McMurtry and Croft 1997).

Amongst the Stigmaeidae, *Agistemus floridanus* was the only species found, with higher population in Bento Gonçalves mainly on Chardonnay cultivar in the same period when *Cal. vitis* was in high population. According to Ferla and Moraes (2003), Eriophyidae are the most appropriate food to *A. floridanus*.

The associated plants evaluated in this study contained 39 species of mites, 17 of them as predators. A higher number was collected in *Plantago tomentosa*, with 163 mites, followed by *Solanum* sp., with 115 mites. On *P. tomentosa* 17 species were observed, on *Rumex* sp., ten species, *Senecio* sp., nine, on *Solanum americanum*, eight, on *Bidens pilosa* and *Richardia brasiliensis*, seven and on *Sida sartaremnensis* six species of mites. On strawberry culture, in Rio Grande do Sul, *B. pilosa*, *Rumex* sp. and *Richardia* sp., also presented higher diversity of predator mites (Ferla *et al.* 2007).

Some associated plants in this study presented low diversity of mites and in 26 vegetal species no mites were found. These results, however, may be under the influence of the sampling frequency whereupon each plant was sampled on different places, so few samples from these

plants have been collected.

Uncultured plants occurred naturally and have spontaneous cultivation growth areas. These, in turn, offer many important requirements to natural enemies, such as host and alternative prey, pollen or nectar, as well as microhabitat not available in monocultures free of associated plants (Van Emden 1965). The greater frequency of these plants in agroecosystems can reduce the pest population, since they can hold large predators diversity, and these can naturally migrate from associated plants to cultivated plants (Altieri *et al.* 2003; Bellini *et al.* 2005).

ACKNOWLEDGMENTS: We thank to Vinícola Miolo and UNIVATES University Center for the financial support and the anonymous referees for suggestions and comments to the manuscript. To thank Samuel Felipe Johann by help in translate the manuscript into English.

LITERATURE CITED

- Altieri, M.A., E.N. Solva and C.I. Nicholls. 2003. *O papel da biodiversidade no manejo de pragas*. Ribeirão Preto: Editora Holos. 226 p.
- Amrine Jr, J.A. 1996. *Keys to the World Genera of the Eriophyoidea*. West Bloomfield: Indira Publishing House. 186 p.
- Amrine Jr, J.A. and T.A. Stasny. 1994. *Catalog of the Eriophyoidea of the world*. West Bloomfield: Indira Publishing House. 798 p.
- Baker, E.W. (1943) 1944. Nuevos tydeidos mexicanos (Acarina). *Revista de la Sociedad Mexicana de Historia Natural* 4(3,4):181-189.
- Baker, E.W. 1968. The genus *Pronematus* Canestrini. *Annals of the Entomological Society of America* 61(5): 1091.
- Baker, E.W. and D.M. Tuttle. 1994. *A guide to the spider mites (Tetranychidae) of the United States*. West Bloomfield: Indira Publishing House. 347 p.
- Baker, E.W., T. Kono, Jr. J.W. Amrine, M. Delfinado-Baker and T.A. Stasny. 1996. *Eriophyid Mites of the United States*. West Bloomfield: Indira Publishing House. 394 p.
- Banks, N., 1904. A Treatise on the Acarina, or Mites. *Proceedings of the United States National Museum*, 28.
- Bellini, M.R., L.G. Chiavegato, E.B. Alves and E.O. Sousa. 2005. Plantas de ocorrência espontânea como substratos alternativos para fitoseídeos (Acari, Phytoseiidae) em cultivos de seringueira *Hevea brasiliensis* Muell. Arg. (Euphorbiaceae). *Revista Brasileira de Zoologia* 22(1): 35-42.
- Berlese, A. 1913. Systema Acarorum genera in families suis disposita. *Acaroteca Italica* 1-2: 3-19.
- Braga, F.M. 1957. Catálogo dos ácaros que vivem nas plantas cultivadas e nos produtos armazenados no Brasil. *Boletim Fitossanitário*, 7(1-2): 31-34.
- Bolland, H.R., J. Gutierrez and C.H.W. Flechtmann. 1998. *World catalogue of the spider mite family (Acari: Tetranychidae)*. Leiden: Brill. 392 p.
- Botton, M., E.R. Hickel and S. de J. Soria. 2003. Pragas; p. 82-107. In T.V.M. Fajardo (ed.). *Uvas para processamento: fitossanidade*. Brasília: Embrapa Informação Tecnológica.
- Botton, M., S. de J. Soria and E.R. Hickel. 2005. Pragas da videira; p.1-22. In E. Hickel, F.F. Calegario, G.W.B. de Melo, G B. Kuhn, J.C. Nachtigal, J.D.G. Maia, J.F. da S. Protas, L.M.R. de Mello, L. da R. Garrido, M.A.F. Conceição, M. Botton, O.R. Sônego, R. de L. Naves, S. de J. Soria, T.V.M. Fajardo and U.A. Camargo (ed.). *Uvas sem sementes, cultivares BRS Morena, BRS Clara e BRS Linda*. Bento Gonçalves: Embrapa Uva e Vinho.
- Chant, D.A. 1959. Phytoseiid mites (Acarina: Phytoseiidae) . Part I. Bionomics of seven species in southeastern England. Part II. A taxonomic review of the family Phytoseiidae, with descriptions of thirty-eight new species. *The Canadian Entomologist Supplement* 12: 1-166.
- Chant, D. A and E.W. Baker. 1965. The Phytoseiidae (Acarina) of Central America. *Memoirs of the Entomological Society of Canada*, Canada, 41, 56 p.
- Chant, D.A. and J.A. McMurtry. 1994. A review of the subfamily Phytoseiinae and Typhlodrominae (Acari: Phytoseiidae). *International Journal of Acarology*, 20(4): 171-228.
- Chant, D.A. and J.A. McMurtry. 2003a. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): part I. Neoseiulini new tribe. *International Journal of Acarology* 29(1): 3-46.
- Chant, D.A. and J.A. McMurtry. 2003b. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): part II. The tribe Kampimodromini Kolodochha. *International Journal of Acarology* 29(3): 179-224.
- Chant, D.A. and J.A. McMurtry. 2004a. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): part III. The tribe Amblyseiini Wainstein, subtribe Amblyseiina N. subtribe. *International Journal of Acarology* 30(3):171-228.
- Chant, D.A. and J.A. McMurtry. 2004b. A review of the subfamily Amblyseiinae muma (Acari: Phytoseiidae) Part. IV Tribe Amblyseiini Wainstein, subtribe Arrenoseiina Chant and McMurtry. *International Journal of Acarology* 30(4): 291-312.
- Chant, D.A. and J.A. McMurtry. 2005a. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): part V. Tribe Amblyseiini, subtribe Proprioseiopsina Chant and McMurtry. *International Journal of Acarology* 31(1): 3-22.
- Chant, D.A. and J.A. McMurtry. 2005b A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): part VI. The tribe Euseiini N. tribe, subtribes Typhlodromalina N. subtribe, Euseiina N. subtribe, and Ricoseiina N. subtribe. *International Journal of Acarology* 31(3):187-224.
- Chant, D.A. and J.A. McMurtry. 2005c. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): part VII. Typhlodromipsini N. tribe. *International Journal of Acarology* 31(4): 315-340.
- Chagas, C.M. 1973. Associação do ácaro *Brevipalpus phoenicis* (Geijskes) à mancha anular do cafeeiro. *O biológico* 39: 229-232.
- Chiavegato, L.G. 1987. A leprose do citros no Estado de São Paulo. *Laranja* 2: 7-18.
- Coll, O.R. de and E.D. Saini. 1992. *Insectos y acaros perjudiciales al cultivo de la yerba mate en la Republica Argentina*. Montecarlo: INTA. 48 p.
- Cooreman, J. 1958. Notes et observations sur les Acariens. VII- *Photia graeca* n. sp. (Acaridia, Canestriniidae) et *Lorryia formosa* n.sp. (Stomatostigmata, Tydeidae). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique Entomologie* 34:1-10.
- Croft, B.A. 1975. *Integrated control of apple mites*. East Lansing: Michigan State University, Cooperative Extension Service (Ext. Bull., E-825). 12 p.
- D'Andréa, G. 1951. *Phyllocoptes vitis*. *Anais do V Congresso Brasileiro de Viticultura e Enologia, Caxias do Sul*. 1-72.
- De Leon, D. 1965. Phytoseiid mites from Puerto Rico with descriptions of new species (Acarina: Mesostigmata). *The Florida Entomologist* 48(2): 121-131.
- De Leon, D. 1966. Phytoseiidae of British Guyana with keys to species (Acarina: Mesostigmata). *Studies on the Fauna of Suriname and other Guyanas*, 8, 81-102.
- De Leon, D. 1967. Some mites of the Caribbean Área. Part I. *Acarina on plants in Trinidad, West Indies*. Lawrence: Alles Press Inc. 66 p.
- Denmark, H.A. and M.H. Muma. 1967. Six new Phytoseiidae from Florida (Acarina: Phytoseiidae). *The Florida Entomologist* 50: 169-180.
- Denmark, H.A. and M.H. Muma. 1973. Phytoseiid of Colombia (Acarina: Phytoseiidae). *Revista Brasileira de Biologia* 33(2): 235-276.
- Donnadieu, A.L. 1875. Recherches pour servir à l'histoire des Tétranyques. *Annales Société Linnéenne Lyons*, Ser. 2 (25): 153-155.
- Duso, C. and E. De Lillo. 1996. Grape; p. 571-582 In E.E. Lindquist, M.W. Sabelis and J. Bruin (ed.). *Eriophyoid mites: their Biology, Natural Enemies and Control*. Amsterdam: Elsevier. 643 p.
- Duso, C., A. Pozzebon, C. Capuzzo, V. Malagnini, S. Otto and M. Borgo. 2004. Grape downy mildew spread and mite seasonal abundance in vineyards: effects on *Tydeus caudatus* and its predators. *Biological Control* 32(1): 143-154.
- Ehara, S. and Moraes, G.J. de 1998. A new species of *Amblyseius* (*Euseius*) (Acari: Phytoseiidae) from citrus in Uruguay. *Entomological Science* 1(1): 59-61.
- El-Banhawy, E. M. 1975. New *Amblyseius* mite from Brazil. *Revista Brasileira de Biologia* 35: 549-552.
- Feres, R.J.F. and G.J. de Moraes. 1998. Phytoseiidae mites (Acari: Phytoseiidae) from woody areas in the State of São Paulo, Brazil. *Systematic and Applied Acarology* 3: 125-132.
- Feres, R.J.F., M.R. Bellini and D.C. Rossa-Feres. 2003. Ocorrência e diversidade de ácaros (Acari, Arachnida) associados a *Tabebuia roseo-alba* (Ridl.) Sand (Biononiaceae), no município de São José do Rio Preto, São Paulo, Brasil. *Revista Brasileira Zoologia* 20(3): 373-378.
- Ferla, N.J. and G.J. de Moraes. 1998. Ácaros predadores em pomares de maçã no Rio Grande do Sul. *Anais da Sociedade Entomológica do Brasil* 27(4): 649-654.
- Ferla, N.J. and G.J. de Moraes. 2002. Ácaros predadores (Acari) em plantas nativas e cultivadas do Estado do Rio Grande do Sul, Brasil. *Revista Brasileira de Zoologia* 19(4): 1011-1031.
- Ferla, N.J. and G.J. de Moraes. 2003. Oviposição de ácaros predadores *Agistemus floridanus* Gonzalez, *Euseius concordis* (Chant) e *Neoseiulus anonymus* (Chant and Baker) (Acari) em resposta a diferentes tipos de alimento. *Revista Brasileira de Zoologia* 20(1): 153-155.
- Ferla, N. J. and G.L. da Silva. 2009. A new species of *Amblyseius* Berlese (Acari, Phytoseiidae) from the state of Rio Grande do Sul, Brazil.

- Revista Brasileira de Entomologia*. 53(4): 509-510.
- Ferla, N.J., G.L. da Silva and G.J. De Moraes. 2010. Description of a new species of *Arrenoseius* Wainstein (Acari: Phytoseiidae) from Brazil and a redescription of a similar species from Argentina. *International Journal of Acarology* 36(1): 15-19.
- Ferla, N.J., M.M. Marchetti and J.C. Siebert. 2005. Acarofauna (Acari) de erva-mate (*Ilex paraguariensis* St. Hil.: Aquifoliaceae) no estado do Rio Grande do Sul. *Biociências* 13(2): 133-142.
- Ferla, N.J., M.M. Marchetti and D. Gonçalves. 2007. Ácaros predadores (Acari) associados à cultura do morango (*Fragaria* sp., Rosaceae) e plantas próximas no Estado do Rio Grande do Sul. *Biota Neotropica* 7(2): 1-8.
- Ferla, N.J. and M. Botton. 2008. Ocorrência do ácaro vermelho europeu associado à cultura da videira no Rio Grande do Sul, Brasil. *Ciência Rural* 38(6): 1758-1761.
- Ferreira, R.C.F., J.V. de Oliveira, F.N.P. Haji and M.G.C. Gondim Jr. 2006. Biologia, exigências térmicas e tabela de vida e fertilidade do ácaro-branco *Polyphagotarsonemus latus* (Banks) (Acari: Tarsonemidae) em videira (*Vitis vinifera* L.) cv. Itália. *Neotropical Entomology* 35(1): 126-132.
- Flaherty, D. L. and M. A. Hoy. 1971. Biological control of Pacific mites and Willamette mites in San Joaquin Valley vineyards: Part. III. Role of tydeid mites. *Researches on Population Ecology* 13:80-96.
- Flechtmann, C.H.W. 1973. *Lorryia formosa* Cooreman, 1958 - um ácaro dos citros pouco conhecido no Brasil. *Ciência e Cultura* 25(12): 1179-1181.
- Flechtmann, C.H.W. 1979. *Ácaros de importância agrícola*. São Paulo: Nobel. 189 p.
- Flechtmann, C.H.W. 1987. On the measurements of *Tydeus formosus* (Cooreman, 1958) (Acari, Prostigmata, Tydeidae). *International Journal of Acarology* 13(3): 217.
- Gondim Jr., M.G.C. and G.J. de Moraes. 2001. Phytoseiid mites (Acari: Phytoseiidae) associated with palm trees (Arecaceae) in Brazil. *Systematic and Applied Acarology* 6: 65-94.
- Garman, P. 1948. Mite species from apple trees in Connecticut. *Connecticut Agricultural Experiment Station* 520: 1-27.
- Geijskes, D.C. 1939. Beitrage zur Kenntniss der europäischen Spinnmilben (Acari: tranychidae), mit besonderer Berücksichtigung der niederländischen Arten. *Mededeelingen Van de Landbouwhoogeschool te Wageningen* 42: 1-68.
- Gonzalez, R.H. 1965. A taxonomic study of the genera *Mediolata*, *Zetzellia* and *Agistemus* (Acari: Stigmaeidae). *University of California Publications in Entomology* 41: 64p.
- Halliday R.B., D.E. Walter and E.E. Lindquist. 1998. Revision of the Australian Ascidae (Acarina: Mesostigmata). *Invertebrate Taxonomy* 12: 1-54.
- Haji, F.N.P., A.N. Moreira, P.R.C. Lopes, R.C.F. Ferreira. J.A. de Alencar and F.R. Barbosa. 2001. Monitoramento e determinação do nível de ação do ácaro-branco na cultura da uva. Petrolina: EMBRAPA Semi-Árido. 7 p.
- Hernades, F.A. and R.J.F. Feres. 2006. Biological cycle of *Lorryia formosa* Cooreman, 1958 (Acari, Tydeidae) on rubber trees leaves: a cause of thelytoky. *Experimental and Applied Acarology* 38: 237-242.
- Hessein, N.A. and T.M. Perring. 1986. Feeding habits of the Tydeidae with evidence of *Homeopronematus anconai* (Acari: Tydeidae) predation on *Aculops lycopersici* (Acari: Eriophyidae). *International Journal of Acarology* 12(4): 215-221.
- Hughes, A. M. 1976. *The mites of stored food and houses*. London: Ministry agriculture, Fishery and Food Technical Bulletin, Second Edition 9. 400 p.
- Jeppson, L.R., H.H. Keifer and E.W. Baker. 1975. *Mites injurious to economic plants*. Berkeley: University of California Press. 614 p.
- Johann, L, C.L. Klock, N.J. Ferla and M. Botton. 2009. Acarofauna (Acari) associada à videira (*Vitis vinifera* L.) no Estado do Rio Grande do Sul. *Biociências* 17(1): 1-19.
- Keifer, H, H. 1977. Eriophyid studies C-13. *Agricultural Research Service*. California: USDA. 24 p.
- Knop, N.F. and M.A. Hoy. 1983. Biology of a Tydeid mite, *Homeopronematus anconai* (n. comb.) (Acari: Tydeidae), important in San Joaquin Valley Vineyards. *Hilgardia* 51(5): 1-30.
- Koch. C.L. 1836. *Deutsche Crustacea, Myriapoda, Arachnida*. Fasc. 1. Regensburg.
- Koch, 1842. *Übersicht Des Arachnidensystems*. 5 Vols. Fasc. 3. Nürnberg: C. H. Zeh.
- Kramer, P. 1877. Grundzüge zur Systematik der Milben. *Archiv für Naturgeschichte* 43: 215-247.
- Krantz, G.W. and D. E. Walter. 2009. *A manual of acarology*. Third Edition. Lubbock: Texas Tech University Press. 509 p.
- Laing, J.E. and N.F. Knop. 1982. Potential use of predaceous mites other than Phytoseiidae for biological control of orchards pests; p. 28-35 In M.A. Hoy, G.L. Cunningham and L. Knutson (ed.). *Biological control of pest by mites*. Berkeley: University of California Press.
- Lindquist, E.E. 1986. The world genera of Tarsonemidae (Acari: Heterostigmata): a morphological, phylogenetical, and systematic revision, with a reclassification family-group taxa in the Heterostigmata. *Memoirs of the Entomological Society of Canada* 136: 1-517.
- Lorenzato, D. 1987. Controle biológico de ácaros fitófagos na cultura da macieira no município de Farroupilha-RS. *Agronomia Sulriograndense* 23(2): 167-183.
- Lorenzato, D. and V. A. Secchi. 1993. Controle biológico de ácaros da macieira no Rio Grande do Sul. *Agronomia Sulriograndense* 20(2): 153-165.
- Lorenzi, H., L. Bacher, M. Lacerda and S. Sartori. 2006. *Frutas Brasileiras e Exóticas Cultivadas (de consumo in natura)*. São Paulo: Instituto Plantarum de Estudos da Flora. 672 p.
- Matioli, A.L., E.A. Ueckermann and C.A.L. Oliveira. 2002. Some stigmaeid and Eupalopsellid mites from citrus orchards in Brazil (Acari: Stigmaeidae and Eupalopsellidae). *International Journal of Acarology* 28(2): 99-120.
- McGregor, E.A. 1914. Four new tetranychids. *Annals of the Entomological Society of America* 7: 354-364.
- McGregor, E.A. 1950. Mites of the family Tetranychidae. *The American Midland Naturalist Journal* 44 (2): 257-420.
- McGregor, E.A. 1954. Two new mites in the genus *Typhlodromus* (Acarina: Phytoseiidae). *Southern California Academy Of Science Bulletin* 53: 89-92.
- McMurtry, J.A., C.B. Hufakker and Van De Vrie. 1970. Ecology of tetranychid mites and their natural enemies: a review. 1. Tetranychid enemies: their biological characters and the impact of spray practices. *Hilgardia* 40(11): 331-390.
- McMurtry, J.A. and B.A. Croft. 1997. Life-styles of phytoseiid mites and their roles in biological control. *Annual Review of Entomology* 42: 291-321.
- Mello, L.M.R. de. 2008. *Vitivinicultura brasileira: Panorama 2007*. Bento Gonzalez: EMBRAPA. 4 p.
- Monteiro, L.B. 1994. Ocorrência de *Polyphagotarsonemus latus* (Banks) (Acari: Tarsonemidae) em videira em Bento Gonçalves, RS, Brasil. *Anais da Sociedade Brasileira de Entomologia* 23(2): 349-350.
- Monteiro, L.B. 2002. Manejo integrado de pragas em macieira no Rio Grande do Sul II. Uso de *Neoseiulus californicus* para o controle de *Panonychus ulmi*. *Revista Brasileira de Fruticultura* 24(2): 395-405.
- Moraes, G.J. de. 2002. Controle biológico de ácaros fitófagos com ácaros predadores; p. 225-237 In J.R. Parra, P.S.M. Botelho, B.S. Corrêa-Ferreira and J.M.S. Bento (ed.). *Controle biológico no Brasil: Parasitóides e predadores*. Barueri: Editora Manole Ltda.
- Moraes, G.J. de and C.H.W. Flechtmann. 2008. *Manual de Acarologia: Acarologia básica e ácaros de plantas cultivadas no Brasil*. Ribeirão Preto: Holos Editora. 308 p.
- Moraes, G.J. de, J.A. McMurtry, H.A. Denmark and C.B. Campos. 2004. A revised catalog of the mite family Phytoseiidae. *Zootaxa* 434: 1-494.
- Moraes, G.J. de and J.A. McMurtry. 1983. Phytoseiid mites (Acarina) of northeastern Brazil, with descriptions of four new species. *International Journal of Acarology* 9 (3): 131-148.
- Moraes, G.J. de, J.A. de Alencar, F. Wenzel Neto and S.M.R. Mergulhão. 1990. Explorations for natural enemies of the cassava green mite in Brazil; p. 351-353 In R.H. Howeler (ed.). *Proceedings of the Eight Symposium of the International Society of Tropical Root Crops*. Bangkok: Thailand.
- Moraes, G.J. de, J.A. McMurtry, H.A. Denmark and C.B. Campos. 2004. A revised catalog of the mite family Phytoseiidae. *Zootaxa* 434: 1-494.
- Nalepa, A. 1898. *Das Tierreich. Eine zusammenstellung und Kennzeichnung der rezenten Tierformen*. Berlin: 4 Lieferung. Acarina. 74 p.
- Nalepa, A. 1905. Neue Gallmilben (27 Fortsetzung). Anzeiger der kaiserlichen Akademie Wissenschaften. *Mathematische-naturwissenschaftliche Klasse* 42 (16): 268.
- Oliveira, C.A.L. de. 1986. Flutuação populacional e medidas de controle do ácaro da leprose *Brevipalpus phoenicis* (Geijskes 1939) em citros. Ver. Téc. Cient. Citricultura. *Laranja* 8: 1-31.
- Oudemans, A. C. 1923. Acarologische Anteekeningen LXXIII. *Entomologische Berichten* 6: 202.
- Oudemans, A. C. 1931 C. Acarologische Anteekeningen CVIII. *Entomologische Berichten* 8 (179): 251-263.
- Oudemans, A. C. 1937. *Kritisch Historisch Overzicht der Acarologie*. Leiden: E. Jedeelt Brill Co. 2758 p.
- Pagenstecher, H.A. 1857. Über Milben, besonders die Gattung *Phytoptus*. *Verhandlungen des Naturhistorisch-Medizinscher Vereins zu Heidelberg* 1: 46-53.
- Penteado, S.R. 1995. Principais pragas da erva-mate e medidas alternativas para o seu controle; p. 109-120 In H. Winge, A.G. Ferreira, J.E.A. Mariath and L.C. Tarasconi (ed.). *Erva-Mate: biologia e cultura no Cone Sul*. Porto Alegre: UFRGS.
- Perrin, T.M. and J.A. McMurtry. 1996. Other predatory arthropods; p. 471-479 In E.E. Lindquist, M.W. Sabelis and J. Bruin. (ed.). *Eriophyoid mites: their biology, natural enemies and control*. Amsterdam: Elsevier.

- 790 p.
- Pritchard, D. W. 1956. A new superfamily of Trombidiform mites with the description of a new family, genus and species (Acarina: Iolinoidea: Iolinidae: Iolina nana). *Annals of the Entomological Society of America* 49 (3): 204-206.
- Rahman, K. A. and A.N. Sapra. 1940. Biology of the mite *Paratetranychus indicus* Hirst, a pest of sugar cane in the Punjab. *Indian Journal of Entomology* 2 (2): 201-212.
- Reis, P.R. and L.A.S. Melo. 1984. Pragas da videira. *Informe Agropecuário* 10 (117): 68-72.
- Roggia, S., J.V.C. Guedes, R.C.R. Kuss, J.A. Arnemann and D. Návia. 2008. Spider mites associated to soybean in Rio Grande do Sul, Brazil. *Pesquisa Agropecuária Brasileira* 43(3): 295-301.
- Schruff, G.A. 1985. Grape; p. 359-366 In W. Helle and M.W. Sabelis (ed.). *Spider mites: Their biology, natural enemies and control*. Amsterdam: Elsevier.
- Smiley, R.L. 1992. *The predatory mite family Cunaxidae (Acari) of the world, with a new classification*. West Bloomfield: Indira Publishing House. 356 p.
- Soria, S. de J., C.H.W. Flechtmann and L.B. Monteiro. 1993. Ocorrência de ácaros brancos ou tropical e outros de impotância agrícola de vinhedos do Rio Grande do Sul, Brasil. In: *Anais do VII Congresso de Viticultura e Enologia, Bento Gonçalves e Garibaldi*, 14-16 de julho de 1993, p. 69-71.
- Sousa, J.S.I. 1996. *Uvas para o Brasil*. Piracicaba: FEALQ. 368 p.
- Strickler, K.N., M. Cushing, M.E. Whalon and B.A. Croft. 1987. Mite (Acari) species composition in Michigan apple orchards. *Environmental Entomologia* 16(1): 30-36.
- Summers F.M. and D.W. Price. 1970. Review of the Mite Family Cheyletidae. *University of California Publications in Entomology* 61: 1-153.
- Thor, S. 1902. On the systematic representation of the Acarinen familien Bdellidae Koch, 1842, Grube, 1856, Eupodidae Koch, 1842 and Cunaxidae Sig Thor, 1902. *Verhandlungen der zoologisch-botanischen Gesellschaft in Wien*, 52: 159-165.
- Van Emden, H.F. 1965. The role of uncultivated land in the biology of crop pests and beneficial insects. *Scientific Horticulture* 17: 121-136.
- Vencato, A., L. Rigon, S. Corrêa, E. Reetz, G.R. Rosa, R.R. Beling. 2007. *Anuário brasileiro da uva e do vinho*. Santa Cruz do Sul: Editora Gazeta Santa Cruz. 128 p.
- Voigts, H. and A. C. Oudemans. 1905. Zur Kenntnis der Milben-Fauna von Bremen. *Abhandlungen herausg. von Naturwissenschaftlichen Vereinen zu Bremen* 18:199-252.
- Zacarias, M.S. and G.J. de Moraes. 2001. Phytoseiid mites (Acari) associated with rubber tress and other euphorbiaceous plants in southeastern Brazil. *Neotropical Entomology* 30(4): 579-586.

RECEIVED: April 2010

LAST REVISED: June 2011

ACCEPTED: July 2011

PUBLISHED ONLINE: July 2011

EDITORIAL RESPONSIBILITY: Ana Lúcia Tourinho